



## Research Article

# STUDY OF GLYCATED HAEMOGLOBIN (HBA1C) LEVEL IN NON DIABETIC IRON DEFICIENCY ANEMIA

<sup>1</sup>\*Dr. Chhabra, R. J., <sup>2</sup>Dr. Rakesh dhadhal and <sup>3</sup>Dr. Kiran Sodvadiya

<sup>1</sup>Department of Biochemistry, PDU govt Medical College and Civil Hospital, Rajkot, Gujarat, India

<sup>2</sup>Department of Pathology, Geetanjali Medical College and Hospital, Udaipur, Rajasthan, India

<sup>3</sup>Department of Biochemistry, Index Medical College and Hospital, Indore, M.P, India

### ARTICLE INFO

#### Article History:

Received 28<sup>th</sup> December, 2014  
Received in revised form  
20<sup>th</sup> January, 2015  
Accepted 26<sup>th</sup> February, 2015  
Published online 31<sup>st</sup> March, 2015

#### Keywords:

Iron Deficiency Anemia,  
HbA1c,  
Hb.

### ABSTRACT

**Background:** Glycated haemoglobin is produced by a ketoamine reaction between glucose and the N-terminal valine of both  $\beta$ -chains of the haemoglobin molecule. The major form of glycated haemoglobin is haemoglobin A1c (HbA1c). The HbA1c fraction is abnormally elevated in chronic hyperglycaemic diabetic patients and it correlates positively with the glycaemic control. However, increased glycated haemoglobin levels have been documented in iron deficiency anaemic patients without any history of diabetes.

**Aims and Objectives:** The aim of this study was to determine the effect of IDA on the HbA1c levels in nondiabetic patients.

**Methods:** 60 non-diabetic, anaemic patients and 60 age-matched healthy subjects were involve in this study. Haematologic investigations were done and the fasting and postprandial glucose and HbA1c levels were measured in all the subjects.

**Results:** The mean HbA1c ( $8.9 \pm 1.8\%$ ) level in the patients with IDA was higher than that in the control group ( $5.4\% \pm 0.8$ ) ( $p < 0.05$ ). There were no differences in the levels of fasting and postprandial glucose between the IDA and the control groups ( $p > 0.05$ ).

**Conclusion:** HbA1c is not affected by the blood sugar levels alone, and there are various confounding factors when HbA1c is measured, especially that of iron deficiency, which is the commonest of the deficiency diseases worldwide. It is hence important to rule out IDA before making a therapeutic decision, based on the HbA1c levels.

Copyright © 2015 Dr. Chhabra et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Protein glycation is a spontaneous reaction that is believed to play a key role in the pathogenesis of many clinical disorders. The glycation of proteins is enhanced by elevated glucose concentrations. The major form of protein glycation with a clinical consideration is glycated haemoglobin (HbA1c). Glycated haemoglobin is produced by a ketoamine reaction between glucose and the N-terminal valine of both  $\beta$ -chains of the haemoglobin molecule. The major form of glycated haemoglobin is haemoglobin A1c (HbA1c) (shekhar *et al.*, 2014). The term HBA1C refers to glycated haemoglobin. It develops when haemoglobin, a protein within red blood cells that carries oxygen throughout your body, joins with glucose in the blood, becoming 'glycated'. By measuring glycated

haemoglobin (HbA1c), clinicians are able to get an overall picture of what our average blood sugar levels have been over a period of weeks/months. For people with diabetes this is important as the higher the HbA1c, the greater the risk of developing diabetes-related complications. The HbA1c fraction is abnormally elevated in patients with chronic hyperglycaemic diabetes mellitus and it correlates positively with the metabolic control (Kim *et al.*, 2010). According to the American Diabetes Association (ADA) guidelines, the value of HbA1c should be kept below 7% in all the diabetics (Mayer *et al.*, 1983). The values which are greater than 7% indicate an increased chance of progression to the dia-betic complications, especially the microvascular ones. When plasma glucose is consistently elevated, the nonenzymatic glycation of haemoglobin increases; this altera-tion reflects the glycaemic history over the previous 2–3 months, since erythrocytes have an average lifespan of 120 days (Shapiro *et al.*, 1979). Diabetes mellitus is a metabolic disease which is caused by absolute or relative insulin

\*Corresponding author: Dr Chhabra, R. J.,  
Department of biochemistry, PDU govt Medical College and civil hospital, Rajkot, Gujarat, India.

deficiency. About 10% of the Indian population suffers from this disease (Mangukiya *et al.*, 2014). HbA1c is majorly affected by the blood glucose levels alone. However, certain studies have proven that the HbA1c levels are altered by various other coexisting factors, along with diabetes, especially that of iron deficiency anaemia, which is a major public health problem in developing countries like India.

According to the World Health Organization (WHO), iron deficiency is the commonest of the deficiency diseases worldwide (Coban *et al.*, 2004). One of the well-studied pathological ill-effects of IDA in the biological system is the glycation of proteins (Lapolla *et al.*, 2005). The nonenzymatic glycation of proteins has pronounced effects on the structure and the function of proteins. The two known factors which can modulate the glycation of proteins are the prevailing concentration of glucose and the half life of the protein (Mayer *et al.*, 1983). But evidences in the literature have documented increased glycosylated protein levels in some non-diabetic pathological states, like iron deficiency anaemia. Some authors have also found that on supplementation with iron therapy, there was a significant decrease in the levels of glycosylated haemoglobin (Chowdhury *et al.*, 1998). Evidence has accumulated, which supports the hypothesis that the glycation reaction, apart from the traditional chronic hyperglycaemia, can be modulated by the iron status of the patient. If the degree of glycation of other proteins in anaemic patients was similar to that of the glycosylated haemoglobin, it would have important clinical implications. Thus, the main aim of the present study was to determine whether the HbA1c levels were increased among the anaemic patients without diabetes. If so, the iron deficiency has to be corrected before any diagnostic or therapeutic decision was made based on the HbA1c level.

## MATERIALS AND METHODS

Blood samples (3ml) were obtained from 60 anaemic patients of the mean age, 45±5 years, among which 30 were males and 30 were females and 60 age-matched healthy subjects. The anaemic patients were recruited from the Medicine Outpatients Department of our institute, PDU medical college and civil hospital, Rajkot, Guajrat, India. The anaemic patients were selected, based on their haemoglobin levels (Hb < 11 g/dl), and on their peripheral blood smears (mostly microcytic hypochromic), which suggested iron deficiency anaemia and on their haematologic investigations (Like MCV, MCH, MCHC) and serum fasting and postprandial glucose levels.

**Exclusion criteria:** The patients who had glucose tolerance abnormalities (impaired glucose tolerance or diabetes mellitus), haemoglobinopathies, haemolytic anaemia, chronic alcohol ingestion, and chronic renal failure were excluded from the study.

### Laboratory investigations

The blood specimens were drawn after an overnight fast. A hematology parameters (haemoglobin (Hb), haematocrit (Hct), mean corpuscular volume (MCV), and mean corpuscular haemoglobin (MCH)); was estimated in Sysmax automated haematology analyser. The HbA1c levels were determined by turbidimetric immunoinhibition. The peripheral blood smears were examined in all the patients.

## RESULTS

All the results are presented as mean ± S.D. The statistical significance of the difference between the groups was evaluated by using online Student's t-test. The correlation was assessed by the partial correlation analysis. A p value of 0.05 was selected as the point of minimal statistical significance. All the parameters which were tested in both the groups have been reported in Table 1. The fasting and the postprandial blood glucose levels confirmed the nondiabetic status. The peripheral blood smears showed a hypochromic microcytic picture. The HbA1c levels were significantly increased among the IDA

**Table 1. Age wise distribution of participant**

Group	Number	Age(year) Mean ± SD
Test(IDA)	60	45±5
Control	60	43±4

**Table 2. Laboratory data of study group**

Parameter	Test	Control
	Group(IDA)(N=60)	Group(N=60)
Hb gm/dl	10.2±1.5	13.1±1.3
PCV %	34.3±4.3	41.4±2.7
MCV fl	79.1±4.9	83.2±4.6
MCH pg	24.5±2.2	32.9±1.7
Fasting blood sugar mg/dl	94.3±9.4	92.8±9.8
Postprandial blood sugar Mg/dl	113.6±6.1	116.5±5.8
HbA1C %	8.9±1.8	5.4±0.8

**Table 3. Comparison of various parameter between Test (n=60) and control (n=60) Group**

Parameter	Group	Number(n)	Result(Mean)	P-value
Hb (gm/dl)	Test	60	10.2	<0.05
	Control	60	13.1	
Fasting blood sugar (mg/dl)	Test	60	94.3	>0.05
	Control	60	92.8	
Postprandial blood sugar (mg/dl)	Test	60	113.6	>0.05
	Control	60	116.5	
HbA1C %	Test	60	8.9	<0.05
	Control	60	5.4	

patients as compared to those in the controls Table 2. The mean HbA1c (8.9 ± 1.8%) level in the patients with IDA was higher than that in the control group (5.4 ± 0.8%) (p < 0.05) Table 3. There were no differences in the levels of fasting and postprandial glucose between the IDA and the control groups (p > 0.05).

## DISCUSSION

Our results suggested that IDA was associated with higher concentrations of HbA1c. Similarly, Brooks *et al.*, (Brooks *et al.*, 1980) showed higher HbA1c concentrations in iron-deficient nondiabetic adults, which decreased to normal after iron replacement. (Hansen *et al.*, 1990) showed normal HbA1c concentrations in iron deficiency, which dropped to subnormal levels after iron supplementation. (Rai *et al.*, 1986) investigated the different methods and no difference was detected among the colorimetric methods, ion-exchange chromatography and affinity chromatography. The commonly performed immunoturbidometric method was performed to determine the HbA1c levels in this study. In anaemic patients, the concentration of glycosylated haemoglobin has been reported to be increased despite the shortened life span of the erythrocytes.

Several mechanisms have been advocated for this increase in the levels of glycated haemoglobin in anaemic patients. It has been proposed that in iron deficiency, the quaternary structure of the haemoglobin molecule may be altered, and that the glycation of the  $\beta$ -globin chains occurs more readily. According to some investigators, the increase in the gly-cated haemoglobin levels in non-diabetic anaemic patients has been mainly attributed to the decrease in the haemoglobin levels in these patients (El-Agouza *et al.*, 2002). But studies which have investigated the glycation levels of other proteins have not been carried out. This study has got a significant relevance because iron deficiency anaemia is very highly prevalent in a tropical country like India. IDA, being a common variable, influences the HbA1c levels when they are estimated by the most commonly employed methods like immunoturbidometry and so, the IDA must be corrected before making any diagnostic or therapeutic decision based on the HbA1c levels. HbA1c is commonly used to assess the long-term blood glucose control in the patients with diabetes mellitus, because the HbA1c value has been shown to predict the risk for the development of many of the chronic complications in diabetes (Stetler C *et al.*, 2000).

### Conclusion

Our results showed that iron deficiency was associated with higher proportions of HbA1c, which could cause problems in the diagnosis of uncontrolled diabetes mellitus in iron-deficient patients. The iron status must be considered during the interpretation of the HbA1c concentrations in Diabetes mellitus. The iron replacement therapy is thus especially important in diabetic patients with iron deficiency, as it would also increase the reliability of the HbA1c determinations.

### REFERENCES

- Brooks, A.P., Metcalfe, J., Day, J.L., Edwards, M.S. 1980. Iron deficiency and glycosylated haemoglobin A1. *Lancet.*; 19(ii):141
- Chowdhury, T.A. and Lasker, S.S. 1998. The elevated gly-cated haemoglobin in nondiabetic patients is associated with an increased mortality in myocardial infarction. *Postgrad Med J.*, 74:480e1
- Coban, E., Ozdogan, M. and Timuragaoglu, A. 2004. The effect of iron deficiency anemia on the levels of haemoglobin A1c in nondiabetic patients. *Acta Haematol*, 112: 126–28.
- El-Agouza, I., Abu Shola, A. and Sirdah, M. 2002. The effect of iron deficiency anaemia on the levels of the haemoglobin subtypes: the possible consequences in a clinical diagnosis. *Clin Lab Haematol*, 24:285e9.
- Hansen P. G., Eriksen, J., Andersen, T. M. and Olesen, L. 1990. Glycosylated haemoglobin (HbA1c) in iron and vitamin B12 deficiency. *J. Intern. Med.*, 227:133–36
- Kim, C., Bullard K. M., Herman, W. H. and Beckles, G. L. 2010. The association between iron deficiency and the HbA1c levels among adults without diabetes in the National Health and Nutrition Examination Survey, 1999–2006. *Diabetes Care.*; 33: 780–85
- Lapolla, A., Traldi, P. and Fedele, D. 2005. The importance of measuring the products of the non enzymatic glycation of proteins. *Clin. Biochem*, 38:103e15.
- Mangukiya, K. and Sharma, N. 2014. Clinical correlation between diabetes mellitus type 2 and VitaminD3 (25 OH-Cholecalciferol) Level, *I.J.S.N.*, VOL.5 (3) 440-442
- Mayer, T. K. and Freedman, Z. R. 1983. Protein glycosylation in diabetes mellitus: a review of the laboratory measurements and of their clinical utilities. *Clin. Chim. Acta.*, 127:147e84
- Mayer, T. K. and Freedman, Z. R. 1983. Protein glycosylation in diabetes mellitus: a review of the laboratory measurements and their clinical utilities. *Clin Chim Acta.*, 127:147e84
- Rai, K. B. and Pattabiraman, T. N. 1986. The glycosylated haemoglobin levels in iron-deficiency anaemia. *Indian J. Med.*, 83:234–36.
- Shapiro, R., McManus, M., Garrick, L., McDonald, M. J. and Bunn H. F. 1979. The nonenzymatic glycolisation of human haemoglobin at multiple sites. *Metabolism.*, 28(suppl):427–30.
- Shekhar, H. K., Mangukiya, A., Kaur, P. and Jadeja. 2014. Effect of iron deficiency anemia on glycation of haemoglobin in nondiabetics, *I.J.S.N.*, VOL.5 (3) 477-79
- Stetler, C., Mueller, B. and Diem, P. 2000. What you always wanted to know about HbA1c. *Schweiz Med. Wochenschr*, 130 :993–1005

\*\*\*\*\*