



REVIEW ARTICLE

SOME CHEMICAL AND PHYSIOLOGICAL PROPERTIES OF NITRIC OXIDE: A MINI REVIEW

Adebola P. Okedere, Towojesusdunjoyinlo D.Odede, Arnold N. Eluagwule and *Raphael C. Mordi

Chemistry Department, College of Science and Technology, Covenant University, Km 10 Idiroko Road, OTA, Ogun State, Nigeria

ARTICLE INFO

Article History:

Received 17th April, 2017
Received in revised form
26th May, 2017
Accepted 14th June, 2017
Published online 26th July, 2017

ABSTRACT

Nitric oxide (NO) is a simple odd-electron molecule that is a primary pollutant. It transmits signals between neurons; playing a role in blood flow and sexual arousal. In this mini review we try to make accessible the properties and functions of this lipophilic molecule to non-specialists.

Keywords:

Nitric oxide, Structure,
Synthesis, Messenger molecule,
Chemical properties,
Physiological properties.

Copyright ©2017, Adebola P. Okedere et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Some Chemistry of NO: Nitric oxide is a colorless, paramagnetic gas that is moderately soluble in water. It is produced in the body to help the millions of cells communicate with one another by transmitting signals throughout the entire body system^[1]. Nitric oxide is a radical that does not dimerize unless cooled to very low temperature at high pressure^[2,3]. Nitric oxide is a stable molecule with an odd number of electrons (11 valence electrons), •N=O; with an unpaired electron in one of the pi (π) antibonding orbital, which means that the molecule is paramagnetic. The bonding between nitrogen and oxygen is intermediate between a double bond and triple bond. The molecular electron configuration for NO is:

[Core electrons](σ_{2s})²(σ*_{2s})²(π_{2p})⁴(σ_{2p})²(π*_{2p})¹. The net bond order is 2.5 and the oxidation state of nitrogen is +2. The molecular orbital description of the bonding in NO is shown below (Figure 1).




Figure 1. Molecular orbital energy level diagram for NO^[4]

*Corresponding author: Raphael C. Mordi,


Chemistry Department, College of Science and Technology, Covenant University, Km 10 Idiroko Road, OTA, Ogun State, Nigeria

Some physical properties of NO are presented in Table 1 below, where it can be seen that its solubility in water is low and heavier than water.

Table 1. Some physical properties of nitric oxide

| Chemical formula | NO |
|---------------------|--|
| Density | 1.3402 g dm ⁻³ |
| Boiling point | -152 °C |
| Melting point | -164 °C |
| Appearance | Colorless gas |
| Molar mass | 30.01 g mol ⁻¹ |
| Solubility in water | 0.0098 g/100 mL (0 °C) 0.0056 g/100 mL (20 °C) |
| Molecular shape | Linear |

Nitric oxide is prepared industrially by the oxidation of ammonia; catalyzed by Pt/Rh catalyst at between 820°C to 950°C and at pressures between 1 atm and 12 atm.



It can be prepared by the reduction of nitrite ions with iodide ions:




The formation of NO from its elements is unfavorable, that is, its enthalpy of formation, ΔH_f° is positive, $\Delta H_f^\circ = +90.2 \text{ kJ/mol}$ ^[2,3]. Nitric oxide, NO, is a primary pollutant, produced in the high temperature combustion cylinders of cars and jet engines



In the atmosphere, it is oxidized to nitrogen dioxide, a brown gas, which is a major contributor to smog.



The NO₂ absorbs ultraviolet light dissociating into NO molecules and O atoms.




Nitric oxide undergoes a series of reactions to produce other chemical compounds and some of the reactions include:

Its reaction with oxygen to give nitrogen dioxide, NO₂, which when added to water gives nitric acid.



It reacts with hydrogen gas to form nitrogen and water



NO reacts with the halogens to form nitrosyl halides, of general formula XNO. For example, nitrosyl chloride, NOCl, this is a very toxic gas^[5,6].




When nitric oxide reacts with transition metals, metal nitrosyls are formed. The metal nitrosyls are highly colored, deep reds, browns, purples or even black. The NO molecule binds to a low oxidation state metal atom. The most common bonding mode of NO is the terminal linear type (M—NO). The angle of the M—N—O varies from 160° to 180°. Here NO is behaving as a 3-electron donor.



Classic examples of this type of bonding are found in the qualitative test for nitrates, "brown-ring" complex $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$; Roussin's red and black salts $\text{K}_2[\text{Fe}_2(\text{NO})_4\text{S}_2]$ and $\text{K}[\text{Fe}_4(\text{NO})_7\text{S}_3]$ obtained by the action of NO on Fe^{II} ion in the presence of S^{2-} ; and sodium nitroprusside, $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]\cdot\text{H}_2\text{O}$.

In the bent $\text{M}-\text{N}-\text{O}$ conformation the NO group behaves as a one electron donor. The bond angle is in the range $120 - 140^\circ$ [7].



NO causes a wide range of health and environmental effects as a result of the various compounds and derivatives of the family of nitrogen oxides. In a series of reactions that occurs in the troposphere as shown in the reaction equations below, NO increases the concentration of $\text{HO}\cdot$ and O_3 . Ozone is formed when NO and volatile organic compounds (VOCs) react in the presence of heat and sunlight. Nitric oxide reacts with stratospheric ozone to form O_2 and nitrogen dioxide [6].



NO acts as catalyst for the conversion of O_3 to O_2 and is therefore considered as a contributor to the depletion of the ozone layer.



Some Physiological Properties of NO

In 1998 Furchgott, Ignarro and Murad won the Nobel Prize for medicine and Physiology for their earlier discovery in 1986 that NO acted as a signaling agent in biological systems. The molecule penetrates membranes and regulates the functions of the cell [8,9,10]. The ability of NO to act as a neurotransmitter arises from its small molecular size and high lipophilicity which allows it to pass through cell walls. It plays a role in a wide range of biological processes such as blood flow and sexual arousal, blood clotting, blood pressure control by dilating arteries; assisting in gastric motility and the immune system's ability to kill tumor cells and intracellular parasites when it is produced in the macrophage, acting as a toxin [11]. NO has been implicated in the mechanism by which nitroglycerin, which has been used in the treatment of angina, works by dilating the blood vessels, thereby decreasing the pressure and increasing blood flow. The brain has been found to be a rich source of NOS and so it has been postulated that NO may be involved in long-term memory.

NO is synthesized *in vivo* from the amino acid L-arginine, which is catalysed by the haem-containing NO synthase (NOS) enzymes. There are three forms of NO synthases: neuronal NOS (nNOS), inducible NOS (iNOS) and endothelial NOS (eNOS). Inducible NOS (iNOS) is a high-output enzyme and its activity does not depend on the presence of Ca^{2+} ions whereas the isoforms nNOS and eNOS depend on Ca^{2+} for their activity. It has been shown that eNOS and nNOS synthesize NO in response to intracellular Ca^{2+} levels and the effectiveness of NOS isoforms depends on their binding with calmodulin [12]. L-arginine is converted to L-citrulline and NO, in a two-step reaction, involving a 5-electron oxidation process, which requires oxygen in addition to the cofactor NADPH.




Figure 2. Synthesis of NO through the conversion of L-arginine to L-citrulline

NO has been linked with some non-human activities, for example, when the blood-sucking insect *Rhodnius prolixus* attacks a victim. Once its saliva is injected into its victim NO causes the blood vessels to expand and inhibiting blood clotting, making the victim an effective blood donor. The victim now releases histamine to aid healing of the wound but this favors the attacking

insect^[4,13]. NO activities have been implicated in plant physiological and biochemical processes, which has led to its classification as a phytohormone^[14,15,16,17]. NO in plants is thought to be generated through enzymatic and non-enzymatic pathways. The enzymatic pathway seems to be catalysed by cytosolic nitrate reductase (cNR), NOS or NOS-like enzymes and nitrite:NO reductase (N:NOR). Non-enzymatic pathway is nitrite dismutation to NO and nitrate at acidic pH values^[18,19,20,21,22]. Even though NOS-like activity has been detected in plants, animal-type NOS has not been found. NO has been found to have significant effect on seed dormancy, growth, senescence, nitrate reductase activity, respiration, stomatal movement, chlorophyll content, photosynthesis and antioxidant system^[21]. NO is also related to the function of the well-known anti-impotence drug Viagra® (*sildenafil*), which works by affecting the concentration of cyclic guanosine monophosphate (cGMP) in the body. On sexual stimulation, NO is released in the body and this leads to the production of cGMP. cGMP triggers relaxation of smooth muscle and an increase in blood flow in the penis, which causes an erection. However an enzyme cGMP phosphodiesterase type 5 (PDE5) destroys cGMP and this causes loss of erection. Viagra® inhibits PDE5, thereby maintaining the concentration of cGMP in the body and duration of erection.




Figure 3. Role of Viagra® in erectile dysfunction; note that both Viagra and cGMP have similar heterocyclic rings^[4]

Conclusion

Nitric oxide has been shown to play a remarkable role in living systems, as it is involved in blood pressure regulation and blood clotting, neurotransmission. With its involvement in this biochemical and physiological processes in both humans and plants, NO, a small inorganic substance deserves the attention it received which led to the award of the Nobel Prize and the attention it continues to receive. However, the mechanism of its action in plants still requires attention as NO-like protein has not been identified.

REFERENCES

- Anderson JF. Structure and Mechanism in salivary proteins from blood-feeding arthropods, *Toxicon* 2010, 56, 1120– 1129 doi.org/10.1016/j.toxicon.2009.11.002
- Besson-Bard A, Pugin A and Wendehenne D. New insights into Nitric oxide signaling in plants, *Ann Rev Plant Biol*, 2008, 59, 21 – 39 DOI:10.1146/annurev.arplant.59.032607.092830
- Chemical and Engineering News: The surprising life of nitric oxide Small, reactive and containing no carbon atoms, nitric oxide is nevertheless emerging as a biochemical of major importance, 1993, 71(51) 26 – 38 DOI: 10.1021/cen-v071no051.p026
- Chemistry³ Introducing Inorganic, Organic and Physical Chemistry, 2nd Edition, Burrow A, Holman J, Parsons A, Pilling G and Price G. Oxford University Press, 2013: Oxford
- Clark J What is nitric oxide and how does it work? (assessed 17 April 2017) <http://www.nutritionexpress.com/showarticle.aspx?articleid=286>
- Furchgott RF. Endothelium-Derived Relaxing Factor: Discovery, Early Studies, and Identification as Nitric Oxide, *Angew. Chem. Int. Ed.* 1999, 38, 1870 – 1880
- Graziano Mand Lamattina L. Nitric oxide and iron in plants: an emerging and converging story, *Trends Plant Sci*, 2005, 10, 4 – 8 doi.org/10.1016/j.tplants.2004.12.004
- Greenwood NN and Earnshaw A. Chemistry of the Elements, Pergamon Press, Oxford, 1989
- Hayat S, Hasan SA, Mori M, Fariduddin Qand Ahmad A. Nitric oxide: Chemistry, biosynthesis and physiological role, in Nitric oxide in plant physiology, Ed. S Hayat, M Mori, J Pichtel and A Ahmad, Wiley-VCH Verlag GmbH & Co, KGaA Weinheim, 2010: pp 1 – 16
- Housecroft CE and Sharpe AG. Inorganic Chemistry, 4th Edition, Pearson: 2012 London

- Ignarro LJ. Nitric Oxide: A Unique Endogenous Signaling Molecule in Vascular Biology, *Angew. Chem. Int. Ed.* 1999, 38, 1882 – 1892
- Jones L and Atkins P. Chemistry Molecules, Matter and Change, 4th Edition, WH Freeman and Company, New York, 1999
- Kotz JC and Treichel P, Jr. Chemistry and Chemical Reactivity, 4th Edition, Saunders Golden Sunburst Series, Saunders College Publishing, Harcourt Brace College Publishers, Forth Worth, 1999
- Lamattina L, Garcia-Mata C, Graziano Mand Pagnussat GC. Nitric oxide: The versatility of an extensive signal molecule, *Annu Rev Plant Biol* 2003, 54 109 – 136 DOI:10.1146/annurev.arplant.54.031902.134752
- Leshem YY. Nitric oxide in plants, in occurrence, Function and Use, 2000, Kluwer Academic Publishers, Dordrecht, The Netherlands
- Murad F. Discovery of Some of the Biological Effects of Nitric Oxide and Its Role in Cell Signaling, *Angew. Chem. Int. Ed.* 1999, 38, 1856 – 1868
- Neill SJ, Desikan R and Hancock JT. Nitric oxide signaling in plants, *New Phytol* 2003, 159, 11 – 35 DOI:10.1046.j.1469-8137.2003.00804.x
- Omer N, Rohilla A, Rohilla S and Kushnoor A. Nitric oxide: role in human biology. *Int J Pharm Sci Drug Res*, 2012, 4(2), 105 – 109.
- Pagnussat GC, Simontacchi M, Puntarulo Sand Lamattina, L. Nitric Oxide Is Required for Root Organogenesis. *Plant Physiology*, 2002, 129, 954 – 956 doi.10.1104/pp.004036
- Stohr C and Ullrich WR. Generation and possible roles of nitric oxide in plant and roots and their apoplastic space, *J Exp Bot* 2002, 53, 2293 – 2303 DOI.org/10.1093.jxb/erf110
- Stohr Cand Stremlau S. Formation and possible roles of nitric oxide in plant roots, *J Exp Bot*, 2006, 57, 463 – 470 doi.org/10.1093/jxb/erj058
- Weller M, Overton T, Burke J, and Armstrong F. Inorganic Chemistry, 6th Edition, *Oxford University Press*: 2010 Oxford
